6. K.K. Jose and Remesh B Nair(2013). Optimum plot size for field trials in Rubber. Paper submitted to J. Indian Soc. Agric. Statist.
7. K.K. Jose and Remesh B Nair (2013) . Field Plot Technique for experiments with rubber. Paper submitted to J.plantn. Crops.
8. Remesh B Nair and K.K.Jose(2013) , Shortening the Breeding cycle in Hevea Brasiliensis. Paper submitted to J. Plant. Crops.

REFERENCE

1. Abeywardena, V.(1970). The efficiency of pre-experimental yield in the calibration of coconut yields. *Ceylon coconut* 0.**21**:85-91.
2. Abraham, S.T.(2001). *Genetic parameters and divergence in certain wild genotypes of Hevea brasiliensis* (Wild.ex Adr.de Juss.) Muell. Arg. Ph.D.Thesis, Mahatma Gandhi University, Kerala, India, 203p.

3. Abraham, T.P., and Vachani, M.V.(1964). Size and shape of plots and blocks in field experiments with transplanted rice crop. *Indian J. agric. Sci.* **35(**2):152-165.

4. Abraham,T.P., Khosala, R.K. , and Agarwal, K.N.(1969). Uniformity trials on black pepper. *Indian J.agric.Sci.,* **39** (8):790-806.

5. Agarwal K.N., Bavappa K.V.A. and Khosala K.N. (1968). A study of size and shape of plots and blocks and optimum number of pre-experimental period on Arecanut. *Indian J. Agri. Sci*., **38**, 444-460.

6. Aidi-daslin: Achmad Bajhaki; Tohar Danakusuma, M and Murdaningsth Haeruman. 1986. Interaksi genotype x lingkungam pada karet dan perannya di dalam seleksi klon (Genotypes x Environment interaction in rubber and their implications in clonal selection). *Bulletin Perkaretan* **4**(1): 23-27.

7. Balakrishnan, R.,Nair,N.V. and Sreenivasan, T.V.(2000). A method for establishing a core collection of *Saccharum officianarum* L.germplasm based on quantitative-morphological data. *Genetic Resources and Crop Evolution*, **47**(1): 1-9.

8. Balfourier, F., Prosperi, J.M., Charmet, G., Goulard, M. and Monestiez, P. (1999). Using spatial patterns of diversity to develop core collections. In: *Core Collections for Today and Tomorrow* (Eds.R.C. Johnson and T. Hodgkin). International Plant Genetic Resources Institute, Italy, pp. 37-48.

9. Bavappa,K.V.A.(1959). Field experimentation in arecanut part I. *Arecanut J*.**10**:116-122.

10. Brown, A.H.D. (1995). The core collection at the cross roads. In: *Core Collections of Plant Genetic Resources*. (Eds. T. Hodgkin, A.H.D. Brown, T.J.L. Hintum and E.A.V Morales). International Plant Genetic Resources Institute, John Wiley & Sons, Chichester, UK, p. 3-19.

11. Brown, A.H.D. and Spillane, C. (1999). Implementing core collections principles, procedures, progress, problems and promise, In: *Core Collections for Today and Tomorrow* (Eds. R.C. Johnson and T. Hodgkin). International Plant Genetic Resources Institute, Rome, Italy, pp. 1-10.

12. Brown, A.H.D.(1989). Core collections: a practical approach to genetic resources management. *Genome,* **31**(2): 818-824.

13. Brown, A.H.D., Grace, J.P. and Speer, S.S. (1987). Designation of a “core” collection of perennial *Glycine. Soybean Genetics Newsletter*, **14**: 59-70.

14. Charmet, G. and Balfourier, F. (1995). The use of geostatistics for sampling a core collection of perennial ryegrass populations. *Genetic Resources and Crop Evolution*, **42**(4): 303-309.

15. Concalves,P.De S.; Cardoso,M.;Santos,I.C.I.; Martins, A.L.M.; Ortolani.A.A. And Colombo.c.a.;1992. Selection of Hevea mother tree adapted to impredictable annual climatic variability. *Braz.J.Genet*. **15**:136-147.

16. Frankel, O.H. (1984). Genetic perspectives of germplasm conservation. In: Genetic manipulation: Impact on Man and Society. (Eds. W.K. Arber, K. Llimensee, W.J. Peacock and P.Starlinger). Cambridge University Press, Cambridge, UK, pp. 161-170.

17. Frankel, O.H. and Brown, A.H.D. (1984). Plant genetic resources today: A critical appraisal. In: *Crop Genetic Resources: Conservation and Evaluation*.

(Eds. J.H.W. Holden and J.T. Williams). George Allen and Unwin, London, UK, pp. 249-257.

18. Galwey, N.W. (1995). Verifying and validating the representativeness of a core collection. In: *Core Collections of Plant Genetic Resources*. (Eds.T. Hodgkin, A.H.D. Brown, T.J.L. Hintum and E.A.V. Morales). International Plant Genetic Resources Institute, John Wiley & Sons, Chichester, UK, pp. 187-198.

19. George M.V., Patten Shetty, M.V. Narasimhayya, G., Nambiar, M.C. and Sathe, K.V. (1983). Optimum size and shape of plots and blocks for cardamom experiments. *J. Plantn. Crops*, **11** : 50 - 59.

20. Concalves, Pde S.; Cardoso, M.; Colombo,C.A.; Ortolani,A.A.; Martins, A.L.M and SANTOS,J.C.I.1990 Variabilidddate genetic da producao annual da sevingueira; estimativas de para metros geneticos esrudo de interacao genotipo x ambiente. *Bragantia,* **49**: 305-320.

21. Concalves,P.de S.; Ondino Cleante Bataglia; Wagner Rodrigues Dos Santosallino Aldo Ortolani; Ivo Segnini JR and Edson HIdeaki shikasho, 1998. Growth in Sao Paulo state, Brazil, *Genetics and Molecular Biology*, **21**(1): 115-122.

22. Govinda Iyer, T.A. (1957). Statistical Analysis of experimental yield data from coconut trees. *Indian Coconut J*. **11**(1) : 106 – 124.

23. Govinda Iyer,T.A.,(1957). Statistical Analysis of experimental yield data from coconut trees. Indian Coconut J. 11(1):106-124.

24. Hamilton, G.J. 1975. Forest mensuration handbook, Forestry Commission Booklet 39, HMSO, London

25. Harch, B.D., Basford, K.E., DeLacy, I.H., Lawrence, P.K. and Cruickshank, A. (1996). Mixed data types and the use of pattern analysis on the Australian groundnut germplasm data. Genetic Resources and Crop Evolution, 43(4): 363-376.

26. Hathey, W.H., and William, E.J., Efficient estimation of the relationship between plot size and variability of crop yields. *Biometrics,* **14**(1):207-222.

27. Hintum, T.J.L.V. (1994). Comparison of marker systems and construction of a core collection in a pedigree of European spring barley. *Theoretical and Applied, Genetics,* **89**(7-8): 991-997.

28. Hintum, T.J.L.V. and Haalman, D. (1994). Pedigree analysis for composing a core collection of modern cultivars, with examples from barley (*Hordeum vulgare s. La*t.). *Theoretical and Applied Genetics*, **88**(1): 70-74.

29. Husch, B., Miller, C.I. and Beers, T.W. 1972. Forest mensuration. The Ronald Press Company, New York

30. Iyer,G.C.,(1968). Influence of plot size on precision in manuring experiments on *Hevea brasilliensis J. Rubb. Res. Instt. Malaya*. **20**:161-172.

31. Iyer.S.S., and Agarwal, K.C.(1970). Optimum size and shape of plots for sugarcane. *Indian J. Agric. Sci*. **40**(2):124-129.

32. Jain, S.K., Qualset, C.O., Bhat, G.M. and Wu, K.K. (1975). Geographical patterns of phenotypic diversity in a world collection of drum wheat. *Crop Science*, **15**(5): 700-704.

33. Jayaraman, K. 2001. A hand book on statistical analysis in forestry research, Kerala Forest Research Institute, Peechi, India.

34. Jayasekera, N.E.M. SamaranayakE, P. and Karunasekera, K.B. 1977. Initial studies on the nature of genotype-environment interactions in some *Hevea cultivars. Journal of Rubber Research Institute of Sri. Lanka*, **54:**33-42.

35. Jayasekera, N.E.M.1984. A review of studies on genotype-environment interactions in *Hevea*. In: *pro.Int.Rubb.Conf.75 years of rubber Research in Sri Lanka, Colombo,* September 17-19..111-116.

36. Kolkarni, G.A., and Abraham, T.P.,(1963). Investigations on the optimum pre-experimental period in field experiments in perennial crops. *J.Indian Soc. Agric. Statist.* **15**(1):175-183.

37. Li, Y.,Cao, Y.S. and Zhung, X. (1996). A phenotype diversity analysis of foxtail millet (*Seturia italic L.P.Beauv*.) Landraces of Chinese origin. *Genetic Resource and Crop Evolution*, **43**(4): 377-384.

38. Li, Y.,Shi,Y.,Cao, Y. and Wang, T. (2004). Establishment of a core collection for maize germplasm preserved in Chinese National Gene bank using geographic distribution and characteritization data. *Genetic Resource and Crop Evolution*, **51**(8): 845-852)

39. Malosetti, M., Abadie, T. and German, S. (2000). Comparing strategies for selecting a core subset for the Uruguayan barley collection. *Plant Genetic Resource Newsletter*, **121**: 20-26.

40. Menon,T.C.M., and Tyagi, B.N., (1971). Optimum size and shape of plots in experiments with mandarin orange. *Indian J.agric. Sci*. 41(1):55-61.

41. Morales, E.A.V., Valois, A.C.C. and Costa, I.R.C. (1995). Core collection for gene banks with limited resources . In: *Core Collections of Plant genetic Resources*. (Eds. T. Hodgkin, A.H.D. Brown, T.J.L. Hintum and E.A.V. Morales). International Plant Genetic Resources Institute, John Wiley & Sons, Chichester, UK, pp. 241-250.

42. Nair, R.B. and Prabhakaran, P.V. (1983). Optimum plot size and shape of plots experiments with cashew. *Agric. Res. J. of Kerala*, **21**(2), 83-86.

43. Nambiar, P.T.N. (1989), Vijayakumar, K.., Nampoothiri, KUK and Abraham, K.J. (1992). Optimum plot size for field trials in oil plam. *J. Plantn. Crops.*, **20**(2), 106-109.

44. Narayanan, R.(1965). Some considerations in deciding the optimum number of recorded trees in a plot for experiments on immature *Heves brasilliensis J. Rubb. Res. Instt. Malaya*. **19**:120-131.

45. Noirot, M., Hamon, and Anthony, F. (1996). The principal component scoring: a new method of constituting a core collection using quantitative data. Genetic Resource and Crop Evolution, 43(1): 1-6.

46. Onokpise, O.U.; Olapade, O and Mekako, H.U. 1986 . Genotype x Environment interactions in *Hevea brasiliensis* (Muell.Arg.*) Indian Journal of Genetics*.(3): 506-514.

47. Pahuja, A.N., and Mehra, R.B., 1981). Plot size for coordinated varietal trials in chick pea. *Indian J. agric. Sci*. **51**(2):80-82.

48. Pankajakshan, A.S.,(1960). A note on the relative contribution of genetic and environmental factors on the yield of uniformity treated coconut trees. *Indian Coconut J*. **14**(2):37-43.

49. Pearce, S.C.(1955). Some considerations in deciding plot size in field experiments with trees and bushes. *J. Indian Soc. Agric. Statist*. **7**(1):23-26.

50. Peeters, J.P. and Martinelli, J.A. (1989). Hierarchical cluster analysis as a tool to manage variation in germplasm collection. Theoretical and Applied Genetics**,** **78**(1):42-48.

51. Peez B.G. and Siller L. (1963). The size of experimental cocao plots. *Cacao Turrilba*, **8** : 11 – 12

52. Peiris, T.S.G. and Thattil, R.O. (1997). Alternative methods to determine plot sizes for tree crops – a case study from coconut data. *Cocos*, **12**, 44-53.

53. Philip, M.S. 1994. Measuring trees and forests, CAB International, U.K. Rubber Board, 2004. *Indian Rubber Statistics*, **27**:1

54. Poligano, G.B., Uggenti, P. and Scippa, G. (2001). Diversity analysis and core collection formation in Bari faba bean germplasm. *Plant Genetic Resource Newsletter*, **125**: 33-38.

55. Poligano, G.B., Alba, E., Uggenti, P. and Scippa, G. (1999). Geographical patterns of variation in Bari faba bean germplasm collection. *Genetics Resource and Crop Evolution*, **46**(2): 183-192.

56. Prabhakaran, P.V., and Thomas, E.J. (1974). Optimum plot size for field experiments with tapioca. *Agric Res. J. Kerala,* **12**(1) : 19 – 23.

57. Prabhakaran, P.V., and Thomas, E.J.(1974). Optimum plot size for field experiments with tapioca. *Agric. Res.J. Kerala,* **12**(1): 19-23.

58. Prabhakaran, P.V., Balakrishnan, S. and Mary George (1978). Optimum plot size for field trials with banana. *Agric. Res. J. Kerala* **16**(1) : 33 – 38.

59. Prabhakaran, P.V., Balakrishnan,S. and Mary George(1978). Optimum plot size for field trials with banana. *Agric.Res. J. Kerala*, **16**(1):33-38.

60. Prabhakaran,P.V.and Nair,R.B.(1983). Efficiency of covariance ananlysis in manorial trials in cashew.*Agric.Res.J. of Kerala*, **21**(2),83-86.

61. Rajeswari Meenattoor J.; Vinod, K.K.; Krishna Kumar. A.K.; SethuraJ,M.R.; Potty, S.N. and Sinha,R.R. 1991. Clone x Environment interaction during early growth phase of *Hevea brasiliensis*. I. Clonal stability on girth. *Indian Journal of Natural Rubber Research*, ***4***(1): 51-54.

62. Rambabu, Agarwal, M.C., Amraj, P.S., and Chinamani, S. (1982). Size and shape of plots and blocks for field trials on natural grass lands in Nilagri hills. *Indian J. Agric. Sci*. **51** (4) : 271 – 275.

63. Sagreiya, K.P. 1967. Forests and Forestry. National Book Trust, New Delhi.

64. Saraswathy, P. and Krishnan,K.S.(1989). Field plot technique for experiments with coconut.*J.Planin.Crops*., 16(Suppl.), 481-487.

65. Saraswathyamma, C.K., Marattukalam, J.G., Chandrasekhar, T.R., Licy, J., Thomas, V., Premakumari, D., Mydin, K.K., Mercykutty, V.C. Meenattoor, J.R., John, A., Gireesh, T., Meenakumari, T., Sankariammal, L., 2000. Rubber clones approved for planting in India -A catalogue. *Rubber Research Institute of India, Kottayam, India*.

66. Sardana, M.G., Sevenath, P.R., and Malhotra, V.P.(1967). Size and shape of plots and blocks in field trials with potato. *Indian J. agric. Sci.* **37**(4) : 338-355.

67. Schoen, D.J. and Brown, A.H.D. (1995). Maximizing genetic diversity in core collections of wild relatives of crop species. In: *Core Collections of Plant Genetic Resources.* (Es. T.Hodgkin, A.H.D. Brown, T.J.L. Hintum and E.A.V. Morales). International Plant Genetic Resources Institute, John Wiley &Sons, Chichester, UK, pp. 55-77.

68. Seena,C. and Prabhakaran, P.V. (1997). Alternative approaches for reduction of error in field experimentation in coconut – an empirical evaluation. *J. Plantn. Crops*., **25**(1), 98-102.

69. Shannon, C.E. and Weaver, W. (1963). *The Mathematical Theory of Communication*. University of Illinois Press, Urbana, IL, USA, 144p.

70. Sheela, M.A. and Unnithan, V.K.G. 1989. Optimum size of plots in cocoa(Theobroma cacao L.) *Plant. Crops* **17** : 18-22.

71. Smith, H.F. 1938. An empirical law describing hotrogenity in the yields of agricultural crops. *J. Agric. Sci*. **28** : 1 – 23.

72. Tan,H. 1991. Genotype x environment (G x E) studies in rubber (Hevea) clones . In: *Proceedings of the 1991 ISOPB International workshop on genotype x environment interaction studies in perennial tree crops*, 12-13 September, 1991, Malaysia. 74-85.

73. Thomas, V., Meenattoor, J.R. and Saraswathyamma, C.K., 2003a. Latex timber clones for small growers in India. Presented at the IRRDB Annual Meeting and Symposium. 15-19 September 2003, Thailand.

74. Thomas, V.,2003b. Rubber wood depot at Perumbavoor, *Rubber*, **447**:8-10.

75. Thompson, S.K., 1992. Sampling. John Wiley & sons., Inc., New York.

76. Tsoumis, G., 1968. Wood as raw material. Pergamon Press. New York.

77. Viswanathan, P. K., George, K.T. and Joseph, T.2003. Timber yield potential of Hevea clones in India: A preliminary assessment. *The Planter, Kuala Lumpur*, **79**:253-261.

78. Viswanathan, P.K., Reghu, C.P. and George, K.T. 2002. From latex to timber: The Indian Perspective. *Rubber Planters' Conference, India*, *200*2, 252-266.

79. Yonezawa, K., Nomura, T. and Morishma, H. (1995). Sampling strategies for use in stratified germplasm collections, In: *Core collections of Plant Genetic Resources* (Eds.T. Hodgkin, A.H.D. Brown, T.J.L. Hintum and E.A.V. Morales). International Plant Genetic Resources Institute, John Wiley & Sons, Chichester,UK, pp. 35-53.

80. Zewdies, Y., Tong, N. and Bosland, P. (2004). Establishing a core collection of Capsicum using a cluster analysis with enlightened selection of accessions. *Genetic Resource and Crop Evolution*, **51**(2): 147-151.